

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**






Fig. 2. Approximate planes for the images of the next figures overlaid on a rendering of the triangulated model for the vertebral surface. The plane for the transverse process images is shown on the left, and the plane for the lamina image is shown on the right.

598405.1 4/20/01




Fig. 3. Actual (top left) and three simulated images (bottom left) of the transverse process with corresponding statistical images (right). The images show only the small, approximately 6 mm axially and 15 mm laterally, portion of the entire image that represents scattering from the transverse process. The statistical images show the variation of the mean, standard deviation and  $SNR_0$  across the image. The simulated images were generated for three different realizations of the microstructure, while the statistical images were computed directly from the model to characterize all possible images.




Fig. 4. Sample simulated images of the lamina and articular processes along with actual image of approximately the same region. From left to right, anatomical structures are the facet joint on the left (Rayleigh scattering with wide axial extent), the lamina in the center (non-Rayleigh scattering with relatively high amplitude) and the inferior articular process on the right (mix of Rayleigh on the sides and non-Rayleigh at the peak).




Fig. 5. Statistical images for the lamina and articular processes. Regions of non-Rayleigh scattering exist along the lamina and the peak of the articular process, although the lamina represents a site of greater  $SNR_0$  than the articular process.




Fig. 6. Samples of a Rayleigh/Gaussian image model, with pixel intensities generated using the computations of previous sections, are shown for the lamina image plane.